



**Clinical Review Criteria
Biofeedback**

NOTICE: Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. (Kaiser Permanente) provide these Clinical Review Criteria for internal use by their members and health care providers. The Clinical Review Criteria only apply to Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. Use of the Clinical Review Criteria or any Kaiser Permanente entity name, logo, trade name, trademark, or service mark for marketing or publicity purposes, including on any website, or in any press release or promotional material, is strictly prohibited.

Kaiser Permanente Clinical Review Criteria are developed to assist in administering plan benefits. These criteria neither offer medical advice nor guarantee coverage. Kaiser Permanente reserves the exclusive right to modify, revoke, suspend or change any or all of these Clinical Review Criteria, at Kaiser Permanente's sole discretion, at any time, with or without notice. **Member contracts differ in health plan benefits. Always consult the patient's Evidence of Coverage or call Kaiser Permanente Member Services at 1-888-901-4636 (TTY 711), Monday through Friday, 8 a.m. to 5 p.m. to determine coverage for a specific medical service.**

Criteria

For Medicare Members

| Source | Policy |
|--|--|
| CMS Coverage Manuals | None |
| National Coverage Determinations (NCD) | Biofeedback Therapy (30.1) . |
| Local Coverage Determinations (LCD) | 3/14/2007 Noridian retired LCD Biofeedback Therapy Policy (L14443) . These services still need to meet medical necessity as outlined in the LCD and will require review. LCDs are retired due to lack of evidence of current problems, or in some cases because the material is addressed by a National Coverage Decision (NCD), a coverage provision in a CMS interpretative manual or an article. Most LCDs are not retired because they are incorrect. Therefore, continue to use LCD L14443 for determining medical necessity. |
| Local Coverage Article | None |

For FEHB plans: See the member's contract for specific coverage details

For Non-Medicare Members

- I. Biofeedback is covered for **ONE of the following**:
 - A. Fecal Incontinence
 - B. Tension or migraine headache if pharmacologic treatment inadequate or not indicated, by **1 or more** of the following:
 - Breast-feeding patient
 - History of long-term, frequent, or excessive use of analgesic or medications that can aggravate headache
 - Insufficient or no response to multiple pharmacologic treatment attempts
 - Intolerance of multiple pharmacologic treatment attempts
 - Patient attempting to become pregnant
 - Pregnant patient

- II. The following indications for biofeedback are not medically necessary. There is insufficient evidence in the published medical literature to show that this service/therapy is as safe as standard services/therapies and/or provides better long-term outcomes than current standard services/therapies.
 - Abdominal pain, recurrent
 - Anxiety disorders
 - Arthritis
 - Asthma
 - Autism
 - Back pain

- Bell's palsy
- Bruxism and sleep bruxism
- Cardiovascular disorders
- Chronic fatigue
- Chronic pain
- Chronic obstructive pulmonary disease (COPD)
- Depression
- Epilepsy
- Facial palsy
- Fibromyalgia
- Hand hemiplegia
- Insomnia
- Knee pain
- Low back pain
- Low vision
- Lupus [systemic lupus erythematosus (SLE)]
- Motor function after stroke, injury, or lower limb surgery
- Movement disorders
- Myalgia or muscle pain
- Neck pain
- Orthostatic hypotension in patients with a spinal cord injury
- Post-traumatic stress disorder (PTSD)
- Raynaud's disease
- Side effects of cancer chemotherapy
- Temporomandibular joint disorders
- Tinnitus
- Vesicoureteral reflux
- Voiding dysfunction
- Vestibulodynia, vulvodynia, vulvar vestibulitis

Biofeedback for the Treatment of Urinary Incontinence

See the [Treatment of Urinary Incontinence criteria document](#)

Neurofeedback for ADHD (EEG Biofeedback)

See the [Neurofeedback criteria document](#)

If requesting this service, please send the following documentation to support medical necessity:

- Last 6 months of clinical notes from requesting provider &/or specialist

The following information was used in the development of this document and is provided as background only. It is provided for historical purposes and does not necessarily reflect the most current published literature. When significant new articles are published that impact treatment option, Kaiser Permanente will review as needed. This information is not to be used as coverage criteria. Please only refer to the criteria listed above for coverage determinations.

Background

Biofeedback is a technique designed to help individuals self-regulate certain physiological processes that are not normally considered to be under voluntary control or responses that are ordinarily easily regulated, but for which regulation has broken down due to trauma or disease. This is achieved through conveying audio and visual information about physiological processes such as blood pressure, heart rate, skin temperature, galvanic skin response (sweating), or muscle tension in real-time to raise awareness of physiological activities and train patients to control them. The goal of biofeedback is that eventually the patient will learn to control physiologic response without the aid of monitors (Kaiser 2011, Roditi 2011).

Different types of biofeedback include (Kaiser 2011, Magnusson 2008, Kapitza 2010):

- Electroencephalography (EEG) biofeedback, which monitors the activity of brain waves linked to different mental states.
- Electrocardiography (EKG) biofeedback, which tracks the patient's heart rate.
- Electromyography (EMG) biofeedback, which uses sensors to measure tension in specific muscles.

- Galvanic skin response biofeedback, which uses sensors to signal anxiety based on the activity of a person's sweat glands and the amount of perspiration on the skin.
- Skin temperature biofeedback, which involves attaching sensors to the fingers or feet to indicate stress when the temperature is low.
- Respiratory biofeedback, which uses sensors to measure breathing.
- Postural biofeedback, which uses sensors to measure body motion.

Biofeedback has been used to treat a variety of medical conditions such as urinary incontinence, ADHD, headaches, anxiety, and back pain.

Evidence and Source Documents

[Biofeedback for Anxiety Disorders](#)

[Biofeedback for Back Pain](#)

[Biofeedback for Migraine and Tension Headaches](#)

[Biofeedback for Treatment of Urinary Incontinence](#)

Medical Technology Assessment Committee (MTAC)

Biofeedback for Anxiety Disorders

02/13/2012: MTAC REVIEW

Evidence Conclusion: There is insufficient evidence to determine the safety and efficacy of biofeedback for the treatment of generalized anxiety disorders.

Articles: The literature search revealed several studies evaluating biofeedback for the treatment of generalized anxiety disorder. All of the studies had small sample sizes and the majority were published more than 20 years ago. The newest study was a randomized controlled trial that evaluated the efficacy of a biofeedback enhanced virtual reality system. This study was not selected for review as the treatment group contained only 4 subjects (Gorini, 2010). Conclusion: There is insufficient evidence to determine the safety and efficacy of biofeedback for the treatment of generalized anxiety disorders.

The use of biofeedback for anxiety disorders does not meet the *Kaiser Permanente Medical Technology Assessment Criteria*.

Biofeedback for Chronic Back Pain

02/13/2012: MTAC REVIEW

Evidence Conclusion: The Kaiser review included four randomized controlled trials that ranged in size from 42 to 128 patients. Findings from these trials suggest that pain and disability improved with biofeedback, cognitive behavioral therapy (CBT), biofeedback plus CBT, placebo biofeedback, and rehabilitation; however, no significant differences were found between biofeedback and the other treatments. The body of evidence was limited by heterogeneity in the patient population, biofeedback protocols, and comparator treatments. Additionally, the studies were small with short follow-up periods. Biofeedback vs. CBT alone vs. waitlisted controls (Newton-John 1995) • N=44 • Type of biofeedback: Electromyography biofeedback (EMG). • Both the biofeedback and the CBT groups showed improvement in pain intensity, pain belief, and depression; however, there no significant differences between the two groups. There was no improvement in the waitlisted control group. Biofeedback plus CBT vs. CBT alone vs. waitlisted controls (Glombiewski 2010) • N=128 • Type of biofeedback: EMG • Both the combined group and the CBT alone group showed improvement in pain intensity compared to waitlisted control; however, there no significant differences between the two groups. Active biofeedback vs. placebo biofeedback (Kapitza 2010) • N=42 • Type of biofeedback: Respiratory biofeedback. • There was no significant difference in pain reduction between the two groups. Biofeedback plus rehabilitation vs. rehabilitation alone (Magnusson 2008) • N=47 • Type of biofeedback: Postural biofeedback. • Although the combined group showed improvements in pain, range of motion, and quality of life, the study did not report if they were statistically significantly different from the rehabilitation alone group. Conclusion: There is insufficient evidence to determine the safety and efficacy of biofeedback for the treatment of chronic back pain.

Articles: The 2007 American College of Physicians and the American Pain Society (ACP/APS) guideline evaluated the evidence on biofeedback for chronic back pain. The studies evaluating this treatment were of poor quality and therefore they were unable to evaluate the net benefits of biofeedback. The conclusions of the ACP/APS guideline were supported by a 2009 BMJ clinical evidence review (Chou 2009). In 2011, the Kaiser Permanente Medical Technology Assessment Team (MTAT) also reviewed biofeedback for the treatment of chronic back pain. No additional studies were identified after the Kaiser review. The following technology assessments were selected for review: Kaiser Permanente TPMG New Medical Technologies. Biofeedback for chronic neck and low back pain. May 2011.

The use of biofeedback for back pain does not meet the *Kaiser Permanente Medical Technology Assessment Criteria*.

Biofeedback for Migraine and Tension Type Headaches

02/13/2012: MTAC Review

Evidence Conclusion: A recent meta-analysis that included 94 RCTs and quasi-experimental studies evaluate the efficacy of different types of biofeedback for the treatment of migraine and tension-type headaches. Results from this analysis suggest that biofeedback was more effective than no treatment for headache reduction in patients with migraine headache (small effect size); however, there was no significant difference between biofeedback and placebo or relaxation. For patients with tension-type headache, biofeedback was significantly more effective than no treatment, placebo, and relaxation for headache reduction (small to medium effect size). There was no significant difference between biofeedback treatment modalities for the reduction of migraine headache pain (Nestouric 2008). A meta-analysis is only as good as the studies that it includes. The studies included in the meta-analysis had several limitations. • The majority of the studies included in the meta-analysis were small. The mean number of subjects per study was 40 for migraine studies and 45 for tension-type headache studies. • The type and number of sessions of biofeedback varied. • Several studies failed to describe basic treatment and patient characteristics. • Several studies used unstructured diagnostic systems. Conclusion: Migraine • Results from a recent meta-analysis suggest that biofeedback may be more effective than no treatment, but not placebo or relaxation for headache reduction. Tension-type headaches • Results from a recent meta-analysis suggest that biofeedback may be more effective than no treatment, placebo, and relaxation for headache reduction. • Another recent BMJ Clinical Evidence review found insufficient evidence to determine whether EMG biofeedback is effective for treating chronic tension-type headaches (Krishnan 2009).

Articles: Several meta-analyses and randomized controlled trials (RCTs) were identified that evaluated the efficacy of biofeedback for the treatment of migraine and tension-type headaches. The most recent meta-analysis was selected for review. An RCT published after the meta-analysis was also identified that evaluated the efficacy of a pain program that included education and training in pain theory plus EMG and temperature biofeedback compared to the pain program alone. This study was not selected for review due to methodological limitations (i.e., small sample size, high loss to follow-up, power not addressed, and baseline characteristics were not presented) (Mullally 2009). The following study was selected for review: Nestoruc Y, Martin A, Rief W, Andrasik F. Biofeedback treatment for headache disorders: a comprehensive efficacy review. *Appl Psychophysiology Biofeedback*. 2008; 33:125-140. See [Evidence Table](#).

The use of biofeedback for Migraine and Tension-type Headaches does not meet the *Kaiser Permanente Medical Technology Assessment Criteria*.

Applicable Codes

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

| CPT® Codes | Description |
|------------|--|
| 90875 | Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (eg, insight oriented, behavior modifying or supportive psychotherapy); 30 minutes |
| 90876 | Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (eg, insight oriented, behavior modifying or supportive psychotherapy); 45 minutes |
| 90901 | Biofeedback training by any modality |
| 90912 | Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; initial 15 minutes of one-on-one physician or other qualified health care professional contact with the patient |
| 90913 | Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; each additional 15 minutes of one-on-one physician or other qualified health care professional contact with the patient (List separately in addition to code for primary procedure) |
| HPCP Codes | Description |
| E0746 | Electromyography (EMG), biofeedback device |

***Note:** Codes may not be all-inclusive. Deleted codes and codes not in effect at the time of service may not be covered.

**To verify authorization requirements for a specific code by plan type, please use the [Pre-authorization Code Check](#).

CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

| Date Created | Date Reviewed | Date Last Revised |
|--------------|--|-------------------|
| 03/06/2012 | 03/06/2012 ^{MDCRPC} , 04/03/2012 ^{MDCRPC} , 02/05/2013 ^{MDCRPC} , 12/03/2013 ^{MPC} , 10/07/2014 ^{MPC} , 08/04/2015 ^{MPC} , 06/07/2016 ^{MPC} , 05/02/2017 ^{MPC} , 02/06/2018 ^{MPC} , 08/07/2018 ^{MPC} , 01/08/2019 ^{MPC} , 01/07/2020 ^{MPC} , 01/05/2021 ^{MPC} , 01/04/2022 ^{MPC} , 01/10/2023 ^{MPC} | 03/12/2024 |

^{MDCRPC} Medical Director Clinical Review and Policy Committee

^{MPC} Medical Policy Committee

| Revision History | Description |
|------------------|---|
| 05/02/2017 | Added indication to cover migraine headaches |
| 07/18/2018 | Added FEHB language |
| 06/23/2020 | Removed deleted CPT code 90911; Added CPT codes 90912 and 90913 |
| 12/02/2022 | Added Retired LCD L14443 |
| 03/12/2024 | Removed urinary conditions from the exclusions list |