



Clinical Review Criteria

Hip Surgery Procedures for Femoroacetabular Impingement Syndrome

NOTICE: Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. (Kaiser Permanente) provide these Clinical Review Criteria for internal use by their members and health care providers. The Clinical Review Criteria only apply to Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. Use of the Clinical Review Criteria or any Kaiser Permanente entity name, logo, trade name, trademark, or service mark for marketing or publicity purposes, including on any website, or in any press release or promotional material, is strictly prohibited.

Kaiser Permanente Clinical Review Criteria are developed to assist in administering plan benefits. These criteria neither offer medical advice nor guarantee coverage. Kaiser Permanente reserves the exclusive right to modify, revoke, suspend or change any or all of these Clinical Review Criteria, at Kaiser Permanente's sole discretion, at any time, with or without notice. **Member contracts differ in health plan benefits. Always consult the patient's Evidence of Coverage or call Kaiser Permanente Member Services at 1-888-901-4636 (TTY 711), Monday through Friday, 8 a.m. to 5 p.m. to determine coverage for a specific medical service.**

Criteria

For Medicare Members

Source	Policy
CMS Coverage Manuals	None
National Coverage Determinations (NCD)	None
Local Coverage Determinations (LCD)	None
Kaiser Permanente Medical Policy	Due to the absence of a NCD, LCD, or other coverage guidance, Kaiser Permanente has chosen to use their own Clinical Review Criteria, " Hip Surgery Procedures for Femoroacetabular Impingement Syndrome " for medical necessity determinations. Use the Non-Medicare criteria below.

For Non-Medicare Members

Service	Criteria
Hip surgery for labral tear repair (29916)	Labral repair (29916) does not require review. When a diagnosis of labral tear has been confirmed with imaging, FAI procedures (29914 and/or 29915) may be covered as part of the repair.
Hip surgery procedures for Femoroacetabular impingement Syndrome (FAI) (29914, 29915)	There is insufficient evidence in the published medical literature to show that this service/therapy is as safe as standard services/therapies and/or provides better long-term outcomes than current standard services/therapies.

If requesting review for these services, please send the following documentation:

- Last 6 months of clinical notes from requesting provider &/or specialist

The following information was used in the development of this document and is provided as background only. It is provided for historical purposes and does not necessarily reflect the most current published literature. When significant new articles are published that impact treatment option, Kaiser Permanente will review as needed. This information is not to be used as coverage criteria. Please only refer to the criteria listed above for coverage determinations.

Background

Femoroacetabular impingement (FAI) syndrome is a recently recognized diagnosis in primarily younger individuals where relatively minor abnormalities in the joint (orientation or morphology) are thought to cause friction/impingement and pain. It is theorized that FAI starts the breakdown of cartilage, leading to osteoarthritis. There are two types of FAI: cam impingement (non-spherical femoral head or abnormality at the head-neck junction) and pincer impingement (deep or retroverted acetabulum resulting in over coverage

of the femoral head). Proponents believe that surgical correction of the impinging deformities will alleviate the symptoms and retard the progression of OA degeneration. Surgery to correct FAI includes arthroscopy, open dislocation of the hip, and arthroscopy combined with a mini-open approach. The purpose of the surgery is to remove abnormal outgrowths of bone and damaged cartilage, and to reshape the femoral neck to ensure that there is sufficient clearance between the rim of the acetabulum and the neck of the femur.

Medical Technology Assessment Committee (MTAC)

Femoroacetabular Impingement Syndrome

06/17/2013: MTAC REVIEW

Evidence Conclusion: There is no new evidence that would change or add to the recommendations of the HTA review as regards the conservative or surgical treatment of femoroacetabular impingement. The results of these non-randomized observational studies as well as other published retrospective series with or without a comparison group should be interpreted with caution. Due to the nature of the study design, they are subject to selection bias, observation bias, confounding and other limitations, and only provide the lowest grade of evidence.

Articles: Larson CM, Giveans R, Stone RM, et al. Arthroscopic debridement versus refixation of the acetabular labrum associated with femoroacetabular impingement. Mean 3.5 –year follow-up. *Am J Sports Med.* 2012; 40:1015-1021. Larson and colleagues (2012) reported on outcomes of two cohorts of patients with femoroacetabular impingement who were treated with either arthroscopic debridement or refixation of the acetabular labrum in one center, but at different time periods. The mean follow-up ranged between 24 and 72 months with a mean of 42 months. The results indicate that the labral fixation was associated with better Harris Hip Scores (HHS), Short Form-12 (SF-12) and visual analog scale (VAS) for pain outcomes compared to arthroscopic focal debridement. Zingg PO, Ulbrich EJ, Buehler TC, et al. Surgical hip dislocation versus hi arthroscopy for femoroacetabular impingement. Clinical and morphological short-term results. *Arch Orthop Trauma Surg.* 2013; 133:69-79. Zingg and colleagues (2013) compared surgical hip dislocation versus hip arthroscopy in 38 patients presenting with clinically FAI that was morphologically verified with plain radiographs and MRI. In 28 of the 38 participants the selection of the procedure was based on the patient’s decision, and only 10 agreed to be randomly allocated to either procedure. There were statistically significant differences in the morphological pathology (in terms of acetabular coverage angle, and head-neck offset ratio) between the two groups at baseline. The primary outcome of the study was the alpha angle on a cross-table view. The results of the study showed that patients in the hip arthroscopy group had faster recovery and better short-term outcomes compared to those treated with surgical hip dislocation. However, the hip arthroscopy showed some overcorrection of the cam deformity and limited frequency of labrum refixations, which the authors indicate that they may lead to negative impact on long-term outcomes.

The use of FIS does not meet the *Kaiser Permanente Medical Technology Assessment Criteria*.

Per the Washington State Health Care Authority Health Technology Clinical Committee (HTCC) coverage determination following Femoroacetabular Impingement Syndrome re-review (adopted 1/17/2020):

Hip surgery for femoroacetabular impingement syndrome is not a covered benefit.

Applicable Codes

Considered not medically necessary:

CPT® Codes	Description
27299	Unlisted procedure, pelvis or hip joint
29914	Arthroscopy, hip, surgical; with femoroplasty (i.e., treatment of cam lesion)
29915	Arthroscopy, hip, surgical; with acetabuloplasty (i.e., treatment of pincer lesion)

***Note:** Codes may not be all-inclusive. Deleted codes and codes not in effect at the time of service may not be covered.

**To verify authorization requirements for a specific code by plan type, please use the [Pre-authorization Code Check](#).

CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Date	Date Reviewed	Date Last
------	---------------	-----------

Created		Revised
08/06/2013	02/04/2013 ^{MPC} , 12/02/2014 ^{MPC} , 10/06/2015 ^{MPC} , 08/02/2016 ^{MPC} , 06/06/2017 ^{MPC} , 04/03/2018 ^{MPC} , 04/02/2019 ^{MPC} , 04/07/2020 ^{MPC} , 04/06/2021 ^{MPC} , 04/05/2022 ^{MPC} , 04/04/2023 ^{MPC}	01/09/2024

^{MPC} Medical Policy Committee

Revision History	Description
06/06/2017	Adopted KP policy for Medicare members
04/07/2020	Removed generic service code 27299 and added more specific codes 29914, 29915 and 29916
04/29/2020	Added CPT codes 27299 and 29862 and ICD-10 codes M25.851, M25.852 and M25.859
04/26/2021	Removed CPT code 29862 and ICD-10 codes M25.851, M25.852 and M25.859
11/06/2021	Removed CPT code 29916
04/05/2022	Added the Washington Health Care Authority HTCC decision from January 2020.
01/09/2024	MPC approved to revise the FAI policy to allow for FAI procedures to be authorized when a separate procedure for labral repair is indicated. 60-day notice is not required.