



**Kaiser Foundation Health Plan
of Washington**

**Clinical Review Criteria
Transanal Endoscopic Resection of Rectal Carcinoma**

NOTICE: Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. (Kaiser Permanente) provide these Clinical Review Criteria for internal use by their members and health care providers. The Clinical Review Criteria only apply to Kaiser Foundation Health Plan of Washington and Kaiser Foundation Health Plan of Washington Options, Inc. Use of the Clinical Review Criteria or any Kaiser Permanente entity name, logo, trade name, trademark, or service mark for marketing or publicity purposes, including on any website, or in any press release or promotional material, is strictly prohibited.

Kaiser Permanente Clinical Review Criteria are developed to assist in administering plan benefits. These criteria neither offer medical advice nor guarantee coverage. Kaiser Permanente reserves the exclusive right to modify, revoke, suspend or change any or all of these Clinical Review Criteria, at Kaiser Permanente's sole discretion, at any time, with or without notice. **Member contracts differ in health plan benefits. Always consult the patient's Evidence of Coverage or call Kaiser Permanente Member Services at 1-888-901-4636 (TTY 711), Monday through Friday, 8 a.m. to 5 p.m. to determine coverage for a specific medical service.**

**Criteria
For Medicare Members**

Source	Policy
CMS Coverage Manuals	None
National Coverage Determinations (NCD)	None
Local Coverage Determinations (LCD)	None
KPWA Medical Policy	Due to the absence of a NCD, LCD, or other coverage guidance, KPWA has chosen to use their own Clinical Review Criteria, " Transanal Endoscopic Resection of Rectal Carcinoma ," for medical necessity determinations. Use the Non-Medicare criteria below.

For Non-Medicare Members

Transanal Endoscopic Microsurgery (TEM) will be considered medically necessary for **ONE** or more of the following indications:

1. Benign rectal tumors (adenomas)
2. Low-risk Tis and T1 rectal carcinoma
3. Small rectal carcinoids (less than 2 cm in diameter)
4. T2 cancer in someone medically unable to undergo a major operation

Kaiser Permanente Washington does not cover Transanal Endoscopic Microsurgery (TEM) for lesions that do not meet the criteria above.

The following information was used in the development of this document and is provided as background only. It is provided for historical purposes and does not necessarily reflect the most current published literature. When significant new articles are published that impact treatment option, Kaiser Permanente will review as needed. This information is not to be used as coverage criteria. Please only refer to the criteria listed above for coverage determinations.

Background

Transanal endoscopic microsurgery (TEM) is a minimally invasive surgical technique that was developed to avoid the morbidity of radical surgery for adenomas and early-stage rectal cancer, while still allowing for complete removal of the lesion. TEM requires specialized instrumentation. TEM uses a natural opening (the anus) to reach the target organ, and is a valuable surgical technique with a low complication rate for patients with appropriate rectal lesions. The main advantages of TEM are preservation of the rectum, anus and fecal continence, low complication rates, short operation times, lower blood loss, shorter hospital stays, and shorter recover times. Other advantages include better exposure, magnified stereoscopic view, and greater reach into the middle and upper rectum.

Local excision (LE) alone does not offer the opportunity for lymph node biopsy and, therefore, has been reserved for patients in whom the likelihood of cancerous extension is small. LE can occur under direct visualization for

rectal tumors within 10 cm of the anal verge and may be most appropriate for small tumors (less than 4cm) confined to the submucosa (T1, as defined by the TNM staging system). TEMS extends local excision ability to the proximal rectosigmoid junction. Adenomas, large rectal polyps (which cannot be removed through a colonoscope), retrorectal masses, small carcinoid tumors, and non-malignant conditions such as strictures or abscesses are amenable to local excision by either method. TEMS can avoid morbidity and mortality associated with major rectal surgery, including the fecal incontinence related to stretching of the anal sphincter, and can be performed under general or regional anesthesia. Use of TEMS for resection of rectal cancers is more controversial.

The most common treatment for rectal cancer is surgery, either open resection or local excision. The technique chosen depends on the size and location of the tumor, evidence of local or distal spread, and patient characteristics and goals. Open, wide resections have the highest cure rate, but may also have significant adverse effects, such as lifelong colostomy, bowel, bladder, or sexual dysfunction. The use of LE in rectal adenocarcinoma is an area of much interest; however, because LE alone does not offer the opportunity for lymph node biopsy it has been reserved for patients in whom the likelihood of cancerous extension is small. Despite this increased risk of local recurrence, local excision may be an informed alternative for patients. TEMS permits local excision beyond the reach of direct visualization equipment.

Applicable Codes

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT® or HCPC Codes	Description
0184T	Excision of rectal tumor, transanal endoscopic microsurgical approach (ie, TEMS), including muscularis propria (ie, full thickness)

***Note:** Codes may not be all-inclusive. Deleted codes and codes not in effect at the time of service may not be covered.

**To verify authorization requirements for a specific code by plan type, please use the [Pre-authorization Code Check](#).

CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions, and materials are copyrighted by Centers for Medicare Services (CMS).

Date Created	Date Reviewed	Date Last Revised
03/07/2017	03/07/2017 ^{MPC} , 05/02/2023 ^{MPC}	

^{MPC} Medical Policy Committee

Revision History	Description
03/07/2017	MPC approved to adopt criteria for TEMS