

1 PATIENT INFORMATION Patient Name: _____ Phone: _____ Address: _____ City: _____ State: _____ Zip: _____ MRN #: _____ DOB: _____ Drug Allergies: _____	2 PRESCRIBER INFORMATION Prescriber's Name: _____ DEA#: _____ NPI: _____ Clinic/Facility Name: _____ Address: _____ City: _____ State: _____ ZIP: _____ Phone: _____ Fax: _____
---	--

3 Instructions to Provider
 All orders with ✓ will be placed unless otherwise noted. Please fax completed order form to 206-326-2139. For drug prior authorization, call 1-888-767-4670 or visit <https://wa-provider.kaiserpermanente.org/provider-manual/clinical-review/preservice>
4 CLINICAL INFORMATION
 Diagnosis (ICD-10 code): _____ Date of Last Dose: _____

5 ABATACEPT PRESCRIPTION INFORMATION
Abatacept (Orencia) Route: Intravenous
 First Dose: No Yes Weight: _____ kg Date Recorded: _____
 Dose: 500 mg 750 mg 1000 mg Other _____
 Initial: Start on week 0 and 2, then continue every 4 weeks thereafter
 Maintenance: Continue every _____ weeks thereafter
 Refills: 11 months Other _____

 Infusion Access: PIV CVAD Other: _____
 Patient's Current Home Care/Specialty Pharmacy: _____

Infusion Reaction Medications & Supplies

<input checked="" type="checkbox"/> Hydrocortisone sodium succinate injectable 100 mg IV Sig: Once PRN for hypersensitivity <input checked="" type="checkbox"/> Epinephrine AutoInjector ? 0.15mg ? 0.3mg QTY: 2 Sig: Inject into lateral thigh muscle for severe allergic reaction. Seek medical attention after use. <input checked="" type="checkbox"/> Sodium Chloride 0.9% IV Flush: Flush 10 ml IV before/after medication administration or as needed for line maintenance. Then flush with 30 ml IV after medication administration.	<input checked="" type="checkbox"/> Diphenhydramine injectable 25 mg IV Sig: Once PRN, may repeat x1 for urticaria, pruritis, shortness of breath <input checked="" type="checkbox"/> Sodium Chloride 0.9% IV 250 mL Bag Sig: Once PRN for anaphylaxis
---	---

 Labs /Special Instructions/Pre-Meds: _____

Infusion Protocol: <ul style="list-style-type: none"> • Infuse per manufacturer guidelines • Monitor vital signs (Temp, BP, HR, RR) every 15 minutes x 4; then every 30 minutes x2; then every 60 minutes until completion of infusion • Documentation must include: <ul style="list-style-type: none"> ○ Start and end time of infusion ○ All rate changes, vital signs, including initial and final set ○ Patient response 	<ul style="list-style-type: none"> • Observe patient for signs of infusion rate-related adverse reactions: <ul style="list-style-type: none"> ○ Blood pressure changes, increased pulse rate ○ Fever, chills ○ Headache ○ Chest, back or hip pain ○ Dyspnea ○ Mild erythema
--	---

6 PHYSICIAN SIGNATURE REQUIRED

X _____ SUBSTITUTION PERMITTED (Date)	X _____ DISPENSE AS WRITTEN (Date)
---	--