

Kaiser Permanente Washington Home Infusion Pharmacy (KPWAHIP) Belatacept (NULOJIX) Prescription Referral Form

	Phone: (206) 326-2990 Fax Referral Io: (206) 326-213
1 PATIENT INFORMATION	2 PRESCRIBER INFORMATION
Patient Na	D 11 / N
Phone:	Prescriber's Name:NPI:
Address:	
City:State:Zip:	Clinic/Facility Name:
MRN #:	Address: State: ZIP:
DOB:	City:State:ZiP:
Drug Allergies:	Phone:Fax:
3 Instructions to Provider	
All orders with ü will be placed unless otherwise noted. Please fax com call 1-888-767-4670 or visit https://wa-provider.kaiserpermanente.org	•
4 CLINICAL INFORMATION	
Diagnosis (ICD-10 code):	Date of Last Dose:
5 BELATACEPT (NULOJIX) PRESCRIPTION INFORMATION	
Nulojix (Belatacept) in 0.9% Sodium Chloride	Route: Intravenous
First Dose: □ No □ Yes	
Weight: kg Date Recorded:	_
Dose & Frequency: mg/kg x weight (kg) once every	_weeks
Refills: □ 11 months □ Other	
Infusion Access: PIV CVAD Other:	
Patient's Current Home Care/Specialty Pharmacy:	
Administer as IV infusion over 30 minutes with 0.2-micron or 1.2-micron filter	
Infusion Reaction Medications & Supplies	
	Diphenhydramine injectable 25 mg IV
Sig: Once PRN for hypersensitivity	Sig: Once PRN, may repeat x1 for urticaria, pruritis,
✓ Epinephrine Auto-Injector □ 0.15mg □ 0.3mg QTY: 2	shortness of breath
	Sodium Chloride 0.9% IV 250ml Bag
reaction. Seek medical attention after use.	Sig: Once PRN for anaphylaxis
✓ Sodium Chloride 0.9% IV Flush: Flush 10 ml IV before/after m	• • • • • • • • • • • • • • • • • • • •
South Chorac 0.5% IV Hash. Hash 10 hir V before, after hi	edication duministration of as needed for line maintenance
Labs /Special Instructions/Pre-Meds:	
Infusion Protocol:	Observe patient for signs of infusion rate-related adverse
Infuse per manufacturer guidelines	reactions:
Monitor vital signs (Temp, BP, HR, RR) every 15 minutes x 4;	 Blood pressure changes, increased pulse rate
then every 30 minutes x2; then every 60 minutes until	o Fever, chills
completion of infusion	Headache
Documentation must include:	Chest, back or hip pain
Start and end time of infusion	Dyspnea
 All rate changes, vital signs, including initial and final set 	Mild erythema
	Educate patients to limit exposure to ultraviolet light and
Patient response	sunlight
6 PHYSICIAN SIGNATURE REQUIRED	
X	x
SUBSTITUTION PERMITTED (Date)	DISPENSE AS WRITTEN (Date)