

<b>1 PATIENT INFORMATION</b> Patient Name: _____ Phone: _____ Address: _____ City: _____ State: _____ Zip: _____ MRN #: _____ DOB: _____ Drug Allergies: _____	<b>2 PRESCRIBER INFORMATION</b> Prescriber's Name: _____ DEA#: _____ NPI: _____ Clinic/Facility Name: _____ Address: _____ City: _____ State: _____ ZIP: _____ Phone: _____ Fax: _____
---	--

**3 Instructions to Provider**

All orders with ✓ will be placed unless otherwise noted. Please fax completed order form to 206-326-2139. For drug prior authorization, call 1-888-767-4670 or visit <https://wa-provider.kaiserpermanente.org/provider-manual/clinical-review/preservice>.

**4 CLINICAL INFORMATION**

Diagnosis (ICD-10 code): \_\_\_\_\_ Date of Last Dose: \_\_\_\_\_

**5 CERAZYME PRESCRIPTION INFORMATION**

**Cerezyme (Imiglucerase)** Route: Intravenous

First Dose:  No  Yes Weight: \_\_\_\_\_ kg Date Recorded: \_\_\_\_\_

Dose:  60 units/kg x weight = \_\_\_\_\_ Unit  Other \_\_\_\_\_ Unit

Frequency: Every 2 weeks or  Other \_\_\_\_\_

Refills: 11 months or  Other \_\_\_\_\_

Infusion Access:  PIV  CVAD  Other: \_\_\_\_\_

Patient's Current Home Care/Specialty Pharmacy: \_\_\_\_\_

**Infusion Reaction Medications & Supplies**

✓ <b>Hydrocortisone sodium succinate injectable 100 mg IV</b> Sig: Once PRN for hypersensitivity	✓ <b>Diphenhydramine injectable 25 mg IV</b> Sig: Once PRN, may repeat x1 for urticaria, pruritis, shortness of breath
✓ <b>Epinephrine Auto-Injector</b> <input type="checkbox"/> 0.15mg <input type="checkbox"/> 0.3mg QTY: 2 Sig: Inject into lateral thigh muscle for severe allergic reaction. Seek medical attention after use.	✓ <b>Sodium Chloride 0.9% IV 250ml Bag</b> Sig: Once PRN for anaphylaxis
✓ <b>Sodium Chloride 0.9% IV Flush:</b> Flush 10 ml IV before/after medication administration or as needed for line maintenance	

**Labs /Special Instructions/Pre-Meds:** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

<p><b>Infusion Protocol:</b></p> <ul style="list-style-type: none"> <li>• Infuse per manufacturer guidelines</li> <li>• Monitor vital signs (Temp, BP, HR, RR) every 15 minutes x 4; then every 30 minutes x2; then every 60 minutes until completion of infusion</li> <li>• Documentation must include:                             <ul style="list-style-type: none"> <li>○ Start and end time of infusion</li> <li>○ All rate changes, vital signs, including initial and final set</li> <li>○ Patient response</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• Observe patient for signs of infusion rate-related adverse reactions:                             <ul style="list-style-type: none"> <li>○ Blood pressure changes, increased pulse rate</li> <li>○ Fever, chills</li> <li>○ Headache</li> <li>○ Chest, back or hip pain</li> <li>○ Dyspnea</li> <li>○ Mild erythema</li> </ul> </li> </ul>
---	---

**6 PHYSICIAN SIGNATURE REQUIRED**

X _____ (Date)	X _____ (Date)
----------------------	----------------------