

## Kaiser Permanente Washington Home Infusion Pharmacy (KPWAHIP) Naglazyme (galsulfase)Prescription Referral Form

Phone: (206) 326-2990 Fax Referral To: (206) 326-2139

1 PATIENT INFORMATION	2 PRESCRIBER INFORMATION
Patient Name:	
Phone:	Prescriber's Name:
Address:	DEA#:NPI:
City:State:Zip:	Clinic/Facility Name:
MRN #:	Address:
DOB:	City:State:ZIP:
Drug Allergies:	Phone:Fax:
3 Instructions to Provider	
All orders with ✓ will be placed unless otherwise noted. Please fax completed order form to 206-326-2139. For drug prior authorization,	
call 1-888-767-4670 or visit https://wa-provider.kaiserpermanente.o	· ·
Table 1 000 7 07 1070 01 Visit Interpretation of the provider interpretation	g, provider mandal, annear review, preservice.
4 CLINICAL INFORMATION	
Diagnosis (ICD-10 code):	Date of Last Dose:
5 NAGLAZYME PRESCRIPTION INFORMATION	
5 NAGLAZYIME PRESCRIPTION INFORMATION	
Naglazyme (Galsulfase)	Route: Intravenous
First Dose: □ No □ Yes	Weight: kg Date Recorded:
Dose: □ 1 mg/kg x weight=mg or □ Other	<del></del>
Frequency: Once weekly or   Other	
Refills:   11 months  Other	
Infusion Access:   PIV CVAD Other:	
Patient's Current Home Care/Specialty Pharmacy:	
Infusion Reaction Medications & Supplies	
,	Diphenhydramine injectable 25 mg IV
Sig: Once PRN for hypersensitivity	Sig: Once PRN, may repeat x1 for urticaria, pruritis,
✓ Epinephrine Auto-Injector □ 0.15mg □ 0.3mg QTY: 2	shortness of breath
	Sodium Chloride 0.9% IV 250ml Bag
reaction. Seek medical attention after use.	Sig: Once PRN for anaphylaxis
✓ Sodium Chloride 0.9% IV Flush: Flush 10 ml IV before/after medication administration or as needed for line maintenance	
Souldin Chioride 6.5% iv Plash. Hash 10 hill iv before, after medication administration of as needed for line maintenance	
Labs /Special Instructions/Pre-Meds:	
Edb3 / Special Histractions/11c Weas.	
Infusion Protocol:	Observe nations for signs of influsion rate related adverse.
<ul> <li>Infuse per manufacturer guidelines</li> </ul>	<ul> <li>Observe patient for signs of infusion rate-related adverse reactions:</li> </ul>
<u> </u>	
Wienter vital signs (remp) by they tree y as minutes x 1,	
then every 30 minutes x2; then every 60 minutes until	<ul><li>Fever, chills</li><li>Headache</li></ul>
completion of infusion	
Documentation must include:	<ul> <li>Chest, back or hip pain</li> </ul>
Start and end time of infusion	O Dyspnea
All rate changes, vital signs, including initial and final set	<ul> <li>Mild erythema</li> </ul>
Patient response	
6 PHYSICIAN SIGNATURE REQUIRED	
X	X
SUBSTITUTION PERMITTED (Date)	DISPENSE AS WRITTEN (Date)