

Kaiser Permanente Washington Home Infusion Pharmacy (KPWAHIP) **Tocilizumab Prescription Referral Form**

	Phone: (206) 326-2990 Fax Referral To: (206) 326-213
1 PATIENT INFORMATION	2 PRESCRIBER INFORMATION
Patient Name:	
Phone:	Prescriber's Name:
Address:	DEA#: NPI:
City:State:Zip:	Clinic/Facility Name:
MRN #:	Address:
DOB:	City:State:ZIP:
Drug Allergies:	Phone:Fax:
3 Instructions to Provider	
All orders with ✓ will be placed unless otherwise noted. Please fax co call 1-888-767-4670 or visit https://wa-provider.kaiserpermanente.org	
4 CLINICAL INFORMATION	
Diagnosis (ICD-10 code):	Date of Last Dose:
5 TOCILIZUMAB PRESCRIPTION INFORMATION	
Tocilizumab-aazg (TYENNE) in 0.9% Sodium Chloride IV Injec	
TB Status: □ PPD (negative) Date: □ □Last Chest	X-Ray: Date Active TB: Unknown
First Dose: □ No □ Yes	
Weight:kg Date Recorded:	
Dose & Frequency: mg/kg x weight (kg) once every	_weeks
Refills: 11 months Other	
Infusion Access: PIV CVAD Other:	_
Patient's Current Home Care/Specialty Pharmacy:	_
•For patients with RA, GCA, COVID-19, CRS, PJIA, and SJIA patients at or abo	ove 30 kg, dilute to 100 mL in 0.9% Sodium Chloride Injection, USP for
intravenous infusion using aseptic technique.	
• For PJIA, SJIA and CRS patients less than 30 kg, dilute to 50 mL in 0.9% Soc	lium Chloride Injection, USP for intravenous infusion using aseptic
technique.	
 Administer as a single intravenous infusion over 60 minutes; do not admin 	nister as bolus or push. Max 800mg.
Infusion Reaction Medications & Supplies	
✓ Hydrocortisone sodium succinate injectable 100 mg IV	Diphenhydramine injectable 25 mg IV
Sig: Once PRN for hypersensitivity	Sig: Once PRN, may repeat x1 for urticaria, pruritis,
✓ Epinephrine Auto-Injector □ 0.15mg □ 0.3mg QTY: 2	shortness of breath
Sig: Inject into lateral thigh muscle for severe allergic	Sodium Chloride 0.9% IV 250 ml Bag
reaction. Seek medical attention after use.	Sig: Once PRN for anaphylaxis
✓ Sodium Chloride 0.9% IV Flush: Flush 10 ml IV before/after m	• • • • • • • • • • • • • • • • • • • •
Labs /Special Instructions/Pre-meds:	
Infusion Protocol:	 Observe patient for signs of infusion rate-related adverse
 Infuse per manufacturer guidelines 	reactions:
 Monitor vital signs (Temp, BP, HR, RR) every 15 minutes x 4; 	 Blood pressure changes, increased pulse rate
then every 30 minutes x2; then every 60 minutes until	 Fever, chills
completion of infusion	o Headache
Documentation must include:	 Chest, back or hip pain
 Start and end time of infusion 	o Dyspnea
 All rate changes, vital signs, including initial and final set 	 Mild erythema
 Patient response 	
6 PHYSICIAN SIGNATURE REQUIRED	
X	х
SUBSTITUTION PERMITTED (Date)	DISPENSE AS WRITTEN (Date)