KAISER PERMANENTE®

Kaiser Permanente Washington Home Infusion Pharmacy (KPWAHIP)
Trastuzumab Prescription Referral Form
Phone: (206) 326-2990 **Fax Referral To: (206) 326-2139**

1 PATIENT INFORMATION

Patient Name: _____
Phone: _____
Address: _____
City:_____**State:**_____**Zip:**_____
MRN #: _____
DOB: _____
Drug Allergies: _____

2 PRESCRIBER INFORMATION

Prescriber's Name: _____
DEA#: _____**NPI:** _____
Clinic/Facility Name: _____
Address: _____
City: _____**State:** _____**ZIP:** _____
Phone: _____**Fax:** _____

3 Instructions to Provider

All orders with ✓ will be placed unless otherwise noted. Please fax completed order form to 206-326-2139. For drug prior authorization, call 1-888-767-4670 or visit https://wa-provider.kaiserpermanente.org/provider-manual/clinical-review/preservice.

4 CLINICAL INFORMATION

Diagnosis (ICD-10 code): _____ **Date of Last Dose:** _____

5 TRASTUZUMAB PRESCRIPTION INFORMATION

Patient will receive trastuzumab-anns (Kanjinti™) unless Herceptin® is otherwise specified: _____
First Dose: □ No □ Yes **Weight:** _____kg **Date Recorded:** _____
Trastuzumab-anns (KANJINTI) Route: Intravenous
✓ Dose: 6 mg/Kg [maintenance] (= ____ mg) in 0.9% Saline

Frequency: Every 21 days until end date of _____
Infusion Access: □ PIV □ CVAD □ Other: _____
Patient's Current Infusion Pharmacy: _____

Infusion Reaction Medications & Supplies

✓ **Hydrocortisone sodium succinate injectable 100 mg IV**
Sig: Once PRN for hypersensitivity

✓ **Epinephrine Auto-Injector** □ 0.15mg □ 0.3mg **QTY**: 2
Sig: Inject into lateral thigh muscle for severe allergic reaction. Seek medical attention after use.

✓ **Diphenhydramine injectable 25 mg IV**
Sig: Once PRN, may repeat x1 for urticaria, pruritis, shortness of breath

✓ **Sodium Chloride 0.9% IV 250 ml Bag**
Sig: Once PRN for anaphylaxis

✓ **Sodium Chloride 0.9% IV Flush**: Flush 10 ml IV before/after medication administration or as needed for line maintenance

Labs /Special Instructions/Pre-Meds: _____
_____
_____

Infusion Protocol:

- Infuse per manufacturer guidelines
- Monitor vital signs (Temp, BP, HR, RR) every 15 minutes x 4; then every 30 minutes x2; then every 60 minutes until completion of infusion
- Documentation must include:
  - Start and end time of infusion
  - All rate changes, vital signs, including initial and final set
  - Patient response
- Observe patient for signs of infusion rate-related adverse reactions:
  - Blood pressure changes, increased pulse rate
  - Fever, chills
  - Headache
  - Chest, back or hip pain
  - Dyspnea
  - Mild erythema

6 PHYSICIAN SIGNATURE REQUIRED

X	X
SUBSTITUTION PERMITTED *(Date)*	DISPENSE AS WRITTEN (Date)

Members with Medicare Part B or D coverage are not required to use this form.