

Infliximab (REMICADE) – Maintenance Infusion Therapy Plan Orders

Page 1 of 3

Name: _____
Kaiser Permanente Member I.D. # _____
Date of Birth _____

Instructions to Provider

Review orders and note any changes. All orders with will be placed unless otherwise noted. Please fax completed order form to the infusion center where the patient will be receiving treatment (see fax numbers at the end of this protocol). Lab orders are not included on this form – place orders via usual method. Lab monitoring is the responsibility of the ordering physician.

Please complete all of the following:

Pre-Service Authorization has been obtained by Kaiser Permanente **Fax:** 1-888-282-2685 **Voice:** 1-800-289-1363

Order Date: _____	Diagnosis: ICD-10 code (REQUIRED): _____
Weight: _____ kg	ICD-10 description _____

General Plan Communication

- Special instructions/notes: _____

Provider Information

- Live vaccines should not be given concurrently.
- Do not combine with tumor necrosis factor (TNF) agents or other biologic DMARDs.

Infusion Therapy

inFLIXimab (REMICADE) in 0.9% sodium chloride IV infusion (250 mL or 500 mL; max concentration 4 mg/ml)

Dose: 3 mg/kg 5 mg/kg _____ mg/kg x weight (kg) = **Total Dose (will be rounded the nearest 100 mg)**

Indicate rounded dose: 300 mg 400 mg 500 mg 600 mg 700 mg _____ mg

Route: Intravenous

Frequency: **Every** ___ **weeks**

Infusion Rate: 10 – 1,100 mL/hr titrated.

250 mL bag: Start infusion rate at 10 mL/hr for 15 minutes, then increase to 20 mL/hr for 15 minutes, then 40 mL/hr for 15 minutes, then 80 mL/hr for 15 minutes, then 150 mL/hr for 15 minutes, then 250 mL/hr until infusion complete.

500 mL bag: Start infusion rate at 20 mL/hr for 15 minutes, then increase to 40 mL/hr for 15 minutes, then 80 mL/hr for 15 minutes, then 160 mL/hr for 15 minutes, then 300 mL/hr for 15 minutes, then 500 mL/hr until infusion complete.

60 min Rapid Infusion Rate: Infuse over 60 min at a rate up to 550 mL/hr, depending on volume of bag. May be infused over 60 min if patient agrees and received at least 4 consecutive infliximab (or biosimilar) infusions over 2 hours with no evidence of infusion reaction. Document qualifications prior to administration of 60 min rapid infusion.

30 min Rapid Infusion Rate: Infuse over 30 min at a rate up to 1,100 mL/hr, depending on volume of bag. May be infused over 30 min if patient agrees and received at least 4 consecutive infliximab (or biosimilar) infusions over 1 hour with no evidence of infusion reaction. Document qualifications prior to administration of 30 min rapid infusion.

If infusion-related reaction:

- 1) STOP infusion immediately; 2) Increase primary infusion to wide open rate; 3) Administer PRN medications per hypersensitivity protocol; 4) Notify MD

Note any changes to above regimen: _____

Provider Signature: _____ **Date:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Infliximab (REMICADE) – Maintenance Infusion Therapy Plan Orders

Page 2 of 3

Name: _____

Kaiser Permanente Member I.D. # _____

Date of Birth _____

Pre-Meds

- acetaminophen (TYLENOL) tablet
Dose: 650 mg Route: Oral Frequency: Once PRN, 30 minutes prior to infliximab infusion (if not taken at home). May also be given once as needed during infusion for achiness, headache, or fever if not given prior to infusion.
- cetirizine (ZYRTEC) tablet
Dose: 10 mg Route: Oral Frequency: Once, at least 30 minutes prior to infliximab infusion (if not taken at home).
- hydrocortisone sodium succinate (SOLU-CORTEF) injectable [not routine; only if breakthrough reaction]
Dose: 50 mg Route: Intravenous Frequency: Once PRN, 30 minutes prior to infliximab infusion in addition to acetaminophen and antihistamine if patient still experiences symptoms with acetaminophen and antihistamine alone.
- Other: _____
Dose: _____ Route: _____ Frequency: Once, 30 minutes prior to infliximab infusion
- No routine pre-medications necessary. Above pre-meds may be given if patient has reaction and requires pre-medications for future doses.

IV Line Care

- 0.9% sodium chloride infusion 250 mL
Rate: 30 mL/hr Route: Intravenous Frequency: Run continuously to keep vein open
Start peripheral IV if no central line

Infusion Reaction Meds

- albuterol (PROVENTIL) nebulizer solution 0.083%
Dose: 2.5 mg Route: Nebulization Frequency: PRN for shortness of breath/wheezing
- diphenhydrAMINE (BENADRYL) injectable
Dose: 25 mg Route: Intravenous Frequency: Once PRN, May repeat x1 for urticaria, pruritis, shortness of breath. May repeat in 15 minutes if symptoms not resolved.
- EPINEPHrine (Epi-Pen) 0.3 mg/0.3 mL IM Auto-Injector
Dose: 0.3 mg Route: Intramuscular Frequency: Once PRN for anaphylaxis. Inject into lateral thigh and hold for 10 seconds. Massage the injected area. Use for patients weighing greater than 27.3 kg (60 lbs). Use amp and 1.5 inch needle for patients with BMI greater than 30. Notify physician if administered.
- MethylPREDNISolone Sod Succ (PF) Inj 125 mg (SOLU-Medrol PF).
Dose : 125 mg Route : IV push Frequency: Once PRN for hypersensitivity reaction. Notify MD if administered.

Lab Review for Nursing

GI/Derm/Rheumatology Indications (when labs available in Epic):

- Ensure CBC, ALT, AST, and Creatinine have been drawn within the last 16 weeks.
- If labs have not been drawn within 16 weeks, proceed with infusion and instruct patient to receive lab draw today.
- If patient has not had labs drawn within 20 weeks, hold infusion and notify provider.

Nursing Orders

- Do not administer infliximab and notify provider if patient has a temperature greater than 100 degrees F, complains of symptoms of acute viral or bacterial illness, or if patient is taking antibiotics for current infection.
- Administer using a low protein-binding 0.2 micron filter.
- Document protocol qualifications in the Therapy Plan if patient will receive rapid infliximab infusion.
- Post infusion observation time per Nursing Standard or as follows: at least 30 minutes for the first 4 infusions, 1-hour for the first rapid 60 min and first rapid 30 min infusion, and per nurse discretion for subsequent infusions.
- Discontinue IV line when therapy complete and patient stabilized.

References

- REMICADE® Prescribing Information. Revised October 2021.

Provider Signature: _____ Date: _____

Printed Name: _____ Phone: _____ Fax: _____

Infliximab (REMICADE) – Maintenance Infusion Therapy Plan Orders

Page 3 of 3

Kaiser Permanente Infusion Locations

Bellevue Medical Center

11511 NE 10th St, Bellevue, WA 98004

Fax: 425-502-3811 Phone: 425-502-3820

Capitol Hill Medical Center

201 16th Ave E, Seattle WA 98112

Fax: 206-326-3624 Phone: 206-326-3180

Everett Medical Center

2930 Maple St, Everett, WA 98201

Fax: 425-261-1578 Phone: 425-261-1566

Olympia Medical Center

700 Lilly Road N.E., Olympia, WA 98506

Fax: 360-923-7609 Phone: 360-923-7600

Riverfront Medical Center – Spokane

W 322 North River Drive, Spokane, WA 99201

Fax: 509-434-3184 Phone: 509-324-6464

Silverdale Medical Center

10452 Silverdale Way NW, Silverdale, WA 98383

Fax: 360-307-7421 Phone: 360-307-7316

Tacoma Medical Center

209 Martin Luther King Jr Way, Tacoma, WA 98405

Fax: 253-596-3351 Phone: 253-596-3350

Provider Signature: _____ Date: _____

Printed Name: _____ Phone: _____ Fax: _____