

Vedolizumab (ENTYVIO) – Maintenance Infusion Therapy Plan Orders

Page 1 of 2

Name: _____

Kaiser Permanente Member I.D. # _____

Date of Birth _____

Instructions to Provider

Review orders and note any changes. All orders with will be placed unless otherwise noted. Please fax completed order form to the infusion center where the patient will be receiving treatment (see fax numbers at the end of this protocol). Lab orders are not included on this form – place orders via usual method. Lab monitoring is the responsibility of the ordering physician.

Please complete all of the following:

Pre-Service Authorization has been obtained by Kaiser Permanente **Fax:** 1-888-282-2685 **Voice:** 1-800-289-1363

Order Date: _____

Diagnosis:

ICD-10 code (REQUIRED): _____

Weight: _____ kg

ICD-10 description _____

General Plan Communication

- Discontinue therapy if no evidence of therapeutic benefit by week 14.
- Special instructions/notes: _____

Provider Information

- Treatment with vedolizumab not recommended in patients with active, severe infections. Consider withholding vedolizumab in patients who develop a severe infection while on treatment.
- Live vaccines should not be given concurrently or within 3 months of discontinuation of therapy.
- Risk of developing Progressive Multifocal Leukoencephalopathy (PML): Monitor for new or worsening neurological signs or symptoms.
- Monitor for signs and symptoms of liver injury, including fatigue, anorexia, right upper abdominal discomfort, dark urine or jaundice.

Infusion Therapy

- Vedolizumab (ENTYVIO) in 0.9% sodium chloride 250 mL IV infusion**

Dose: 300 mg

Route: Intravenous

Frequency: **Every 8 weeks**

Infusion Rate: Infuse over 30 minutes, starting 60 minutes after treatment start time

If infusion-related reaction:

- 1) STOP infusion immediately;
- 2) Increase primary infusion to wide open rate;
- 3) Administer PRN medications per hypersensitivity protocol;
- 4) Notify MD

Note any changes to above regimen: _____

Pre-Meds

- acetaminophen (TYLENOL) tablet
Dose: 650 mg *Route:* Oral *Frequency:* Once, 30 minutes prior to vedolizumab infusion.
May also be given once as needed during infusion for achiness, headache, or fever.
- cetirizine (ZYRTEC) tablet
Dose: 10 mg *Route:* Oral
Frequency: Once, at least 30 minutes prior to vedolizumab infusion (if not taken at home).
- hydrocortisone sodium succinate (SOLU-CORTEF) injectable
Dose: 50 mg *Route:* Intravenous *Frequency:* Once PRN, 30 minutes prior to vedolizumab infusion in addition to acetaminophen and antihistamine if patient still experiences symptoms with acetaminophen and antihistamine alone.
- Other: _____
Dose: _____ *Route:* Oral *Frequency:* Once, 30 minutes prior to vedolizumab infusion
- No routine pre-medications necessary. Above pre-meds may be given if patient has reaction and requires pre-medications for future doses.

Provider Signature: _____ **Date:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Vedolizumab (ENTYVIO) – Maintenance Infusion Therapy Plan Orders

Page 2 of 2

Name: _____

Kaiser Permanente Member I.D. # _____

Date of Birth _____

IV Line Care

- 0.9% sodium chloride infusion 250 mL
Rate: 30 mL/hr Route: Intravenous Frequency: Run continuously to keep vein open
Start peripheral IV if no central line
- heparin flush 100 unit/mL
Dose: 500 units Route: Intracatheter Frequency: PRN for IV line care per Nursing Policy

Infusion Reaction Meds

- albuterol (ACCUNEB) nebulizer solution
Dose: 2.5 mg Route: Nebulization Frequency: PRN for shortness of breath/wheezing
- diphenhydramine (BENADRYL) injectable
Dose: 25 mg Route: Intravenous Frequency: Once PRN, May repeat x1 for urticaria, pruritus, shortness of breath. May repeat in 15 minutes if symptoms not resolved.
- EPINEPHrine (EpiPen) 0.3 mg/0.3 mL IM Auto-Injector
Dose: 0.3 mg Route: Intramuscular Frequency: Once PRN for anaphylaxis. Inject into lateral thigh and hold for 10 seconds. Massage the injected area. Use for patients weighing greater than 27.3 kg (60 lbs). Use amp and 1.5 inch needle for patients with BMI greater than 30. Notify physician if administered.
- hydrocortisone sodium succinate (SOLU-CORTEF) injectable
Dose: 100 mg Route: Intravenous Frequency: Once PRN for hypersensitivity

Lab Review for Nursing

- Ensure CBC, AST and ALT have been drawn within the last 8 weeks.
- If labs have not been drawn within 8 weeks, proceed with infusion and instruct patient to receive lab draw today.
- Notify provider if patient is more than 12 weeks overdue for labs.

Nursing Orders

- Monitor patient for hypersensitivity reaction: urticaria, dizziness, fever, rash, rigors, pruritus, nausea, flushing, hypotension, dyspnea, and/or chest pain.
- Monitor for signs and symptoms of infection such as fever (greater than 100 degrees F), chills, pruritis, sore throat, erythema, skins sores, and dysuria.
- Discontinue IV line when therapy complete and patient stabilized.

Kaiser Permanente Infusion Locations

Bellevue Medical Center

11511 NE 10th St, Bellevue, WA 98004

Fax: 425-502-3512 Phone: 425-502-3510

Capitol Hill Medical Center

201 16th Ave E, Seattle WA 98112

Fax: 206-326-2104 Phone: 206-326-3109

Everett Medical Center

2930 Maple St, Everett, WA 98201

Fax: 425-261-1578 Phone: 425-261-1566

Olympia Medical Center

700 Lilly Road N.E., Olympia, WA 98506

Fax: 360-923-7106 Phone: 360-923-7164

Riverfront Medical Center – Spokane

W 322 North River Drive, Spokane, WA 99201

Fax: 509-324-7168 Phone: 509-241-2073

Silverdale Medical Center

10452 Silverdale Way NW, Silverdale, WA 98383

Fax: 360-307-7421 Phone: 360-307-7316

Tacoma Medical Center

209 Martin Luther King Jr Way, Tacoma, WA 98405

Fax: 253-383-6262 Phone: 253-596-3666

Provider Signature: _____ Date: _____

Printed Name: _____ Phone: _____ Fax: _____